

DOCKET NO.: AREN-0315

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:

Dominic P. Behan et al.

Serial No.:

Group Art Unit: unknown

Filing Date: January 23, 2002

Examiner: unknown

**For: NON-ENDOGENEOUS, CONSTITUTIVELY ACTIVATED
HUMAN SEROTONIN RECEPTORS AND SMALL
MOLECULE MODULATORS THEREOF**

Assistant Commissioner for Patents
Washington, D.C. 20231

PRELIMINARY AMENDMENT

Please make the following amendments to the above-identified application.

In the Sequence Listing:

Please delete the Sequence Listing on file and insert therefore pages 1-19 comprising the most recently filed Sequence Listing.

In the Specification:


Please amend the paragraph on page 31, lines 2 to 7, to read as follows:

Based upon these results, structure activity analysis of the 103487 compound suggested that a series of derivatives of N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][(4-trifluoromethoxy)phenyl] aminocarboxamide would exhibit similar 5-HT_{2A} activity and selectivity. A series of derivatives of N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][(4-trifluoromethoxy)phenyl] aminocarboxamide were synthesized. These "directed" library


compounds (Tripos, Inc.) were then analyzed in accordance with the protocols of Examples 9c(1), 9c(2) and 9d.

Please amend the paragraph on page 31, lines 8 to 12, to read as follows:


This series of compounds exhibits highly selective 5-HT_{2A} activity. Accordingly, in the first aspect of the invention, a series of compounds exhibiting 5-HT_{2A} receptor activity that are useful as inverse agonists at such receptors is designated by the general formula (A):




Please amend the paragraph on page 34, line 1, to read as follows:




Please amend the paragraph on page 36, lines 9-10, to read as follows:




Please amend the paragraph on page 37, lines 3-4, to read as follows:



Please amend the paragraph on page 37, lines 8-9, to read as follows:



Please amend the paragraph on page 39, lines 1-2, to read as follows:




Please amend the paragraphs spanning pages 39 to page 41 to read as follows:

Compound No.	R ¹	R ²	R ³	R ⁴	X	U	IP ₃ % of Control	IP ₃ AP-3 IC ₅₀ nM	WT 5HT _{2A} LSD IC ₅₀ nM
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][(4-methylthiophenyl)amino]carboxamide									
116079	SCH ₃	H	H	H	O	NH	16	17	4
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][(4-chlorophenyl)amino]carboxamide									
116081	Cl	H	H	H	O	NH	10	3.2	11
{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(4-fluorophenyl)carboxamide									
116082	F	H	H	H	O	NH	11	-	7
{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-[2-(trifluoromethoxy)phenyl]carboxamide									

116087	H	H	CF ₃ O	H	O	NH	11	-	200
{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(2-nitrophenyl)carboxamide									
116089	H	H	NO ₂	H	O	NH	27	-	238
{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(4-methoxyphenyl)carboxamide									
116091	MeO	H	H	H	O	NH	12	-	19
{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(2-methylphenyl)carboxamide									
116092	H	H	Me	H	O	NH	32	-	131
{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-[4-(trifluoromethyl)phenyl]carboxamide									
116097	CF ₃	H	H	H	O	NH	11	-	65
{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(3-chlorophenyl)carboxamide									
116105	H	Cl	H	H	O	NH	11	-	39
{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-[2-chlorophenyl]carboxamide									
116108	H	H	Cl	H	O	NH	6	-	249
{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-[4-(methylethyl)phenyl]carboxamide									
116110	isopropyl	H	H	H	O	NH	7	-	338
{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(3-methoxyphenyl)carboxamide									
116111	H	MeO	H	H	O	NH	7	-	106

[[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino]-N-(3-methylphenyl)carboxamide									
116112	H	Me	H	H	O	NH	14	-	57
[[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino]-N-methyl-N-[4-(trifluoromethoxy)phenyl]carboxamide									
116113	CF ₃ O	H	H	H	O	NCH ₃	-	193	2
N-[4-(tert-butyl)phenyl][3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}carboxamide									
116119	t-butyl	H	H	H	O	NH	17	-	476
N-[4-(dimethylamino)phenyl][3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}carboxamide									
116122	NMe ₂	H	H	H	O	NH	9	-	309
N-(3,5-dichloro-4-methylphenyl)[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}carboxamide									
116138	Me	Cl	H	Cl	O	NH	23	-	122
{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-[4-(trifluoromethylthio)phenyl]carboxamide									
116139	CF ₃ S	H	H	H	O	NH	12	-	56
{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(2-fluorophenyl)carboxamide									
116144	H	H	F	H	O	NH	12	-	37


2-([3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino)carbonylamino)benzamide									
116145	H	H	CONH ₂	H	O	NH	31	-	7473
{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(4-cyanophenyl)carboxamide									
116147	CN	H	H	H	O	NH	12	-	2
{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(2-cyanophenyl)carboxamide									
116148	H	H	CN	H	O	NH	30	-	348



Please amend the table on page 42 to read as follows:

Compound No.		IP ₃ AP-3 IC ₅₀ nM	WT 5HT _{2A} LSD IC ₅₀ nM
116141	N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-[cyclohexylamino]carboxamide	114	81

Please amend the paragraph on page 42, lines 3-4, to read as follows:




Please amend the table on page 42, lower, to 43, upper, to read as follows:

Compound No.	R ¹	R ²	R ³	R ⁴	R ⁵	IP ₃ AP-3	WT 5HT _{2A}
--------------	----------------	----------------	----------------	----------------	----------------	----------------------	----------------------


						IC ₅₀ nM	LSD IC ₅₀ nM
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-[phenylmethylamino]carboxamide							
116143	H	H	H	H	H	120	47
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-[{(4-fluorophenyl)methyl}amino]carboxamide							
116182	F	H	H	H	H	89	132
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-[{(3,4-dimethoxyphenyl)methyl}amino]-carboxamide							
116183	OMe	OMe	H	H	H	-	1010
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-[{(3,4,5-trimethoxyphenyl)methyl}amino]-carboxamide							
116184	OMe	OMe	H	OMe	H	-	2960
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][{(2-methylphenyl)methyl}amino]carboxamide							
116185	H	H	Me	H	H	-	769
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][{(4-methoxyphenyl)methyl}amino]carboxamide							
116189	OMe	H	H	H	H	-	102

Please amend the paragraphs on pages 43-44, to read as follows:



Compound No.	R ¹	R ²	R ³	R ⁴	R ⁵	IP ₃ AP-3	WT 5HT _{2A} LSD
						IC ₅₀ nM	IC ₅₀ nM
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][{2-(4-methoxyphenyl)ethyl} amino]carboxamide							
116194	OMe	H	H	H	H	32	61


Please amend the paragraph on page 44, lines 7-8, to read as follows:



Please amend the paragraph on page 44, line 17, to page 45, line 2, to read as follows:

116100


N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][4-methoxyphenoxy]carboxamide




Please amend the paragraph on page 45, lines 6-7 to read as follows:

116192

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(1,1-dimethylethoxy)carboxamide



Please amend the paragraph on page 47, lines 3-4 to read as follows:




Please amend page 47, line 13, to page 49, line 1 to read as follows:

116101

$$m = 0, R^1 = H, R^4 = 4\text{-trifluoromethoxyphenyl}$$


N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][4-trifluoromethoxyphenyl]carboxamide



116102

m=0, R¹= H, R⁴= thiophene


N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][2-thienyl]carboxamide




116120

m=0, R¹= H, R⁴= chlorophenyl

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][4-chloro-phenyl]carboxamide



Please amend page 51, line 6, to page 52, line 1 to read as follows:




Name	Compound No.	R ¹	R ²	R ³	R ⁴	IP ₃	LSD
						IC ₅₀ nM	IC ₅₀ nM
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-[4-(trifluoromethoxy)phenyl]acetamide	116137	OCF ₃	H	H	H	-	106
N-[3-(4-bromo-2-methylpyrazol-3-	116174	H	F	H	H	153	318

yl)phenyl]-2-(3-fluorophenyl)acetamide							
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(3-methoxyphenyl)acetamide	116175	H	OMe	H	H	108	625
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(2-fluorophenyl)acetamide	116176	H	H	F	H	129	662
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(4-nitrophenyl)acetamide	116177	NO ₂	H	H	H	61	108
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(2-methoxyphenyl)acetamide	116178	H	H	OMe	H	165	2300


compound names not provided

Please amend the paragraph on page 52, line 7 to read as follows:




Please amend the paragraph on page 55, lines 5-13, to read as follows:


Compounds of general formula (I) can be obtained *via* a variety of synthetic routes all of which would be familiar to one skilled in the art. The reaction of isocyanates with amines is a commonly practiced method for the formation of ureas (see Org. Syn. Coll. Vol. V, (1973), 555). Amine (IV), 3-(4-bromo-2-methylpyrazole-3-yl)phenylamine, commercially available from Maybridge Chemical Company, Catalog No. KM01978, CAS No. 175201-77-1] reacts readily with isocyanates (V) in inert solvents such as halocarbons to yield the desired ureas of general formula (I) wherein $R^1 = R^2 = H$:




(IV)



(V)




(I)



Please amend the paragraph spanning page 55, line 14, to page 56, line 1, to read as follows:


Alternatively the amine (IV) can be converted to the corresponding isocyanate

(VI) by the action of phosgene or a suitable phosgene equivalent, *e.g.* triphosgene, in an inert solvent such as a halocarbon in the presence of an organic base such as triethylamine or ethyldiisopropylamine. Isocyanate (VI) reacts with amines of general formula (VII), in an analogous fashion to that described above for the reaction of (IV) with (V), yielding the desired ureas of general formula (I) wherein $R^1 = H$:




Please amend the paragraph spanning page 56, line 2, to page 57, line 1, to read as follows:

Alternatively wherein the isocyanate of general formula (V) is not commercially available it can be prepared from the corresponding amine of general formula (VIII) in an analogous procedure to that described above for the preparation of (VI). Reaction of these isocyanates with (IV) would again yield the requisite ureas of general formula (I) wherein $R^1 = R^2 = H$:



Please amend the paragraph spanning page 57, line 2, to page 58, line 1, to read as follows:


Amines of general formula (VII) are also readily converted to activated isocyanate equivalents of general formula (IX) by the sequential action of carbonyldiimidazole and methyl iodide in tetrahydrofuran and acetonitrile respectively (R.A. Batey *et al*, *Tetrahedron Lett.*, (1998), 39, 6267-6270.) Reaction of (IX) with (IV) in an inert solvent such as a halocarbon would yield the requisite ureas of general formula (I) wherein $R^1 = H$:



(VII)



(IX)

(IV)



(I) R¹ = H

Please amend the paragraph spanning page 58, line 2, to page 59, line 1, to read as follows:


Amine (IV) may be monomethylated according to the procedure of J. Barluenga *et al*, *J. Chem. Soc., Chem. Commun.*, (1984), 20, 1334-1335, or alkylated according to the procedure of P. Marchini *et al*, *J. Org. Chem.*, (1975), 40(23), 3453-3456, to yield compounds of general formula (X) wherein R¹ = lower alkyl. These materials may be reacted as above with reagents of general formula (V) and (IX) as depicted below:




Please amend the paragraph on page 60, line 3, to read as follows:




Please amend the paragraph on page 60, line 4, to read as follows:

(X) R¹ = lower alkyl




(XI)

(II) R¹ = lower alkyl


Please amend the paragraph on page 61, line 1, to read as follows:





Please amend the paragraph on page 62, line 1, to read as follows:



Please amend the paragraph on page 61, line 2, to read as follows:




(X) R¹ = lower alkyl






(III) R¹ = lower alkyl


Please amend page 63, line 1 to read as follows:



(IV)

(III) $R^1 = H$

Please amend page 63, line 5, to read as follows:



Please amend the paragraph on page 66, line 28 to page 67, line 2, to read as follows:

Experiment 2

Preparation and Analysis of 116100

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][4-methoxyphenoxy]carboxamide

To 4-methoxyphenylchloroformate (19 mg, 0.10 mmol) in CH_2Cl_2 (0.5 mL) was added dropwise a solution 3-(3-aminophenyl)-4-bromo-2-methylpyrazole (25 mg, 0.10 mmol) and triethylamine (14 μL , 0.10 mmol) in CH_2Cl_2 (0.5 mL). The mixture was stirred for 16 h and concentrated. Chromatography on flash silica (40% EtOAc/hexane) gave the title compound as a colourless solid (21 mg, 52%), m.p. 140.3-141.8°C. (EtOAc/hexane).

Please amend the paragraph on page 67, lines 9-17, to read as follows:

Experiment 3

Preparation and Analysis of 116101

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][4-trifluoromethoxyphenyl]carboxamide

To 4-(trifluoromethoxy)benzoyl chloride (19 μL , 0.12 mmol) in CH_2Cl_2 (1 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole (30 mg, 0.12 mmol) and triethylamine (17 μL , 0.12 mmol) in CH_2Cl_2 (0.5 mL). The reaction mixture was stirred for 16 h and concentrated. Chromatography on flash silica (50% EtOAc/hexane) gave the title compound as a colourless solid (40 mg, 76%), m.p. 138.6-139.6 °C (EtOAc/hexane).

Please amend the paragraph on page 67, lines 23-30, to read as follows:

Experiment 4

Preparation and Analysis of 116102

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][2-thienyl]carboxamide

To thiophene-2-carbonyl chloride (11 μL , 0.09 mmol) in CH_2Cl_2 (1 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole (25 mg, 0.09 mmol) and triethylamine

(14 μ L, 0.09 mmol) in CH_2Cl_2 (0.5 mL). The reaction mixture was stirred for 16 h and concentrated. Chromatography on flash silica (50% EtOAc/hexane) gave the title compound as a colourless solid (24 mg, 68%), m.p. 127.8-128.6 $^\circ\text{C}$ (EtOAc/hexane).

Please amend the paragraph on page 68, lines 5-14, to read as follows:

Experiment 5

Preparation and Analysis of 116115

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][{(4-trifluoromethoxy)phenyl)methyl}amino]
carboxamide

To a stirred solution of triphosgene (12 mg, 0.04 mmol) in CH_2Cl_2 (0.5 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole (30 mg, 0.12 mmol) and triethylamine (33 μ L, 0.24 mmol) in CH_2Cl_2 (0.5 mL). After 1 h, 4-(trifluoromethoxy)benzylamine (23 mg, 0.12 mmol) was added. The reaction mixture was stirred for 16 h and concentrated. Chromatography on flash silica (75% EtOAc/hexane) gave the title compound as a colourless solid (38 mg, 68%), m.p. 144.6-145.8 $^\circ\text{C}$ (EtOAc/hexane).

Please amend the paragraph on page 68, lines 21-28, to read as follows:

Experiment 6

Preparation and Analysis of 116120

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][4-chlorophenyl]carboxamide

To 4-chlorobenzoyl chloride (15 mg, 0.08 mmol) in CH_2Cl_2 (1 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole (21 mg, 0.08 mmol) and triethylamine (12 μ L, 0.08 mmol) in CH_2Cl_2 (0.5 mL). The mixture was stirred for 16 h and concentrated. Chromatography on flash silica (50% EtOAc/hexane) gave the title compound as a colourless solid (23 mg, 72%), m.p. 184.4-184.8 $^\circ\text{C}$ (EtOAc/hexane).

Please amend the paragraph on page 69, lines 1-33, to read as follows:

Experiment 7

Preparation and Analysis of 116137

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-[4-(trifluoromethoxy)phenyl]acetamide

A solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole (35 mg, 0.14 mmol) and triethylamine (23 μ L, 0.17 mmol) in DMF (0.5 mL) was added in one portion to a stirred solution of 4-trifluoromethoxyphenylacetic acid (31 mg, 0.14 mmol), HBTU (53 mg, 0.14 mmol) and HOBT (19 mg, 0.14 mmol) in DMF (1 mL). The mixture was heated at 70 °C for 24 h and then quenched with aqueous sodium bicarbonate solution. Ethyl acetate was added and the organic phase separated, washed with water (.times.3), brine, dried (MgSO_4) and evaporated. Chromatography on flash silica (50% EtOAc/hexane) gave the title compound as a colourless solid (43 mg, 68%). m.p. 141.2-142.5 °C (EtOAc/hexane).

Please amend the paragraph on page 69, lines 21-33, to read as follows:

Experiment 8

Preparation and Analysis of 116174

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(3-fluorophenyl)acetamide

A mixture of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole (30 mg, 0.12 mmol), 3-fluorophenylacetic acid (18 mg, 0.12 mmol), 1-hydroxybenzotriazole hydrate (16 mg, 0.12 mmol) and 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluoro-phosphate (46 mg, 0.12 mmol) were dissolved in chloroform (1.5 ml). N,N-Diisopropylethylamine (0.02 ml, 0.13 mmol) was added and the mixture stirred at room temperature for 16 h. The reaction mixture was then poured into brine and the organic layer washed with further brine, dried over magnesium sulphate and then concentrated in vacuo. The crude product was purified by column chromatography (ethyl acetate-toluene, 1:1), giving the title compound (12 mg, 26%). Rf 0.41 (ethyl acetate-toluene, 1:1).

Please amend the paragraph on page 70, lines 5-16, as follows:

Experiment 9

Preparation and Analysis of 116175

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(3-methoxyphenyl)acetamide

A solution of 3-methoxyphenylacetyl chloride (0.02 ml, 0.12 mmol) in dichloromethane (0.75 ml) was added dropwise at 0 °C to a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole (30 mg, 0.12 mmol) and triethylamine (0.02 ml, 0.13 mmol) in dichloromethane (0.75 ml). The resulting mixture was stirred at room temperature for 16 h and then poured into brine. The organic layer was washed with more brine then dried over magnesium sulphate and concentrated in vacuo. The crude product was purified by column chromatography (ethyl acetate-toluene, 1:1), giving the title compound (9 mg, 19%). Rf 0.30 (ethyl acetate-toluene, 1:1).

Please amend the paragraph on page 71, lines 21-33, to read as follows:

Experiment 10

Preparation and Analysis of 116176

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(2-fluorophenyl)acetamide

A mixture of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole (30 mg, 0.12 mmol), 2-fluorophenylacetic acid (18 mg, 0.12 mmol), 1-hydroxybenzotriazole hydrate (16 mg, 0.12 mmol) and 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluoro-phosphate (46 mg, 0.12 mmol) were dissolved in chloroform (1.5 ml). N,N-Diisopropylethylamine (0.02 ml, 0.13 mmol) was added and the mixture stirred at room temperature for 16 h. The reaction mixture was then poured into brine and the organic layer washed with further brine, dried over magnesium sulphate and then concentrated in vacuo. The crude product was purified by column chromatography (ethyl acetate-toluene, 1:1), giving the title compound (15 mg, 32%). Rf 0.52 (ethyl acetate-toluene, 1:1).

Please amend the paragraph on page 71, lines 5-17, to read as follows:

Experiment 11

Preparation and Analysis of 116177

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(4-nitrophenyl)acetamide

A mixture of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole (30 mg, 0.12 mmol), 4-nitrophenylacetic acid (22 mg, 0.12 mmol), 1-hydroxybenzotriazole hydrate (16 mg, 0.12 mmol) and 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (46 mg, 0.12 mmol) were dissolved in chloroform (1.5 ml). N,N-Diisopropylethylamine (0.02 ml, 0.13 mmol) was added and the mixture stirred at room temperature for 16 h. The reaction mixture was then poured into brine and the organic layer washed with further brine, dried over magnesium sulphate and then concentrated in vacuo. The crude product was purified by column chromatography (ethyl acetate-toluene, 1:1), giving the title compound (9 mg, 18%). Rf 0.19 (ethyl acetate-toluene, 1:1).

Please amend the paragraph on page 71, line 22, to page 72, line 2, to read as follows:

Experiment 12

Preparation and Analysis of 116178

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(2-methoxyphenyl)acetamide

A mixture of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole (30 mg, 0.12 mmol), 2-methoxyphenylacetic acid (20 mg, 0.12 mmol), 1-hydroxybenzotriazole hydrate (16 mg, 0.12 mmol) and 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (46 mg, 0.12 mmol) were dissolved in chloroform (1.5 ml). N,N-Diisopropylethylamine (0.02 ml, 0.13 mmol) was added and the mixture stirred at room temperature for 16 h. The reaction mixture was then poured into brine and the organic layer washed with further brine, dried over magnesium sulphate and then concentrated in vacuo. The crude product was purified by column

chromatography (chloroform-methanol, 99:1), giving the title compound (18 mg, 38%) as a colourless solid. Rf 0.65 (chloroform-methanol, 98:2).

Please amend the paragraph on page 72, lines 8-15, to read as follows:

Experiment 13

Preparation and Analysis of 116192

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(1,1-dimethylethoxy)carboxamide

To di-tert-butyl dicarbonate (36 mg, 0.17 mmol) in methanol (1 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole (42 mg, 0.17 mmol) in methanol (1 mL). The mixture was stirred for 16 h and concentrated. Chromatography on flash silica (40% EtOAc/hexane) gave the title compound as a colourless solid (29 mg, 49%) (EtOAc/hexane).

Please amend the paragraph on page 72, lines 22 to page 73, line 2, to read as follows:

One or the other (as indicated) of the two following synthetic protocols was used to generate each of the compounds below:

Protocol A:

To an isocyanate (1 mmol) in CH_2Cl_2 (4 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole (1 mmol) in CH_2Cl_2 (4 mL). The mixture was stirred for 16 hours and concentrated. Chromatography on flash silica (20%-80% EtOAc/hexane) followed by recrystallisation gave the pure urea.

Protocol B:

To a stirred solution of triphosgene (0.33 mmol) in CH_2Cl_2 (4 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole (1 mmol) and triethylamine (2 mmol) in CH_2Cl_2 (4 mL). After 1 hour, an aniline was added (1 mmol). The reaction mixture was stirred

for 16 hours and concentrated. Chromatography on flash silica (20%-80% EtOAc/hexane) followed by recrystallisation gave the pure urea.

Please amend the paragraph on page 73, lines 5-8, to read as follows:

Experiment 14

Preparation and Analysis of 116079

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][(4-methylthiophenyl)amino]carboxamide

Please amend the paragraph on page 73, lines 16-18, to read as follows:

Experiment 15

Preparation and Analysis of 116081

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][(4-chlorophenyl)amino]carboxamide

Please amend the paragraph on page 73, lines 29-31, to read as follows:

Experiment 16

Preparation and Analysis of 116082

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(4-fluorophenyl)carboxamide

Please amend the paragraph on page 74, lines 6-8, to read as follows:

Experiment 17

Preparation and Analysis of 116087

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-[2-(trifluoromethoxy)phenyl]carboxamide

Please amend the paragraph on page 74, lines 17-19, to read as follows:

Experiment 18

Preparation and Analysis of 116089

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(2-nitrophenyl)carboxamide

Please amend the paragraph on page 74, lines 29-31, to read as follows:

Experiment 19

Preparation and Analysis of 116091

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(4-methoxyphenyl)carboxamide

Please amend the paragraph on page 75, lines 7-9, to read as follows:

Experiment 20

Preparation and Analysis of 116092

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(2-methylphenyl)carboxamide

Please amend the paragraph on page 75, lines 18-20, to read as follows:

Experiment 21

Preparation and Analysis of 116097

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-[4-(trifluoromethyl)phenyl]carboxamide

Please amend the paragraph on page 75, lines 28-30, to read as follows:

Experiment 22

Preparation and Analysis of 116105

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(3-chlorophenyl)carboxamide

Please amend the paragraph on page 76, lines 8-10, to read as follows:

Experiment 23

Preparation and Analysis of 116108

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(2-chlorophenyl)carboxamide

Please amend the paragraph on page 76, lines 20-22, to read as follows:

Experiment 24

Preparation and Analysis of 116110

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-[4-(methylethyl)phenyl]carboxamide

Please amend the paragraph on page 76, lines 31-33, to read as follows:

Experiment 25

Preparation and Analysis of 116111

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(3-methoxyphenyl)carboxamide

Please amend the paragraph on page 77, lines 8-10, to read as follows:

Experiment 26

Preparation and Analysis of 116112

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(3-methylphenyl)carboxamide

Please amend the paragraph on page 77, lines 19-22, to read as follows:

Experiment 27

Preparation and Analysis of 116113

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-methyl-N-[4-(trifluoromethoxy)phenyl]-carboxamide

Please amend the paragraph on page 77, lines 30-32, to read as follows:

Experiment 28

Preparation and Analysis of 116119

N-[4-(tert-butyl)phenyl]{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}carboxamide

Please amend the paragraph on page 78, lines 8-10, to read as follows:

Experiment 29

Preparation and Analysis of 116122

N-[4-(dimethylamino)phenyl]{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}carboxamide

Please amend the paragraph on page 78, lines 19-21, to read as follows:

Experiment 30

Preparation and Analysis of 116138

N-(3,5-dichloro-4-methylphenyl){[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}carboxamide

Please amend the paragraph on page 78, lines 30-32, to read as follows:

Experiment 31

Preparation and Analysis of 116139

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-[4-(trifluoromethylthio)phenyl]carboxamide

Please amend the paragraph on page 79, lines 8-10, to read as follows:

Experiment 32

Preparation and Analysis of 116141

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(cyclohexyl)carboxamide

Please amend the paragraph on page 79, lines 21-23, to read as follows:

Experiment 33

Preparation and Analysis of 116143

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(phenylmethyl)carboxamide

Please amend the paragraph on page 80, lines 1-3, to read as follows:

Experiment 34

Preparation and Analysis of 116144

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(2-fluorophenyl)carboxamide

Please amend the paragraph on page 80, lines 11-13, to read as follows:

Experiment 35

Preparation and Analysis of 116145

2-({[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}carbonylamino)benzamide

Please amend the paragraph on page 80, lines 21-23, to read as follows:

Experiment 36

Preparation and Analysis of 116147

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(4-cyanophenyl)carboxamide

Please amend the paragraph on page 80, lines 31-33, to read as follows:

Experiment 37

Preparation and Analysis of 116148

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(2-cyanophenyl)carboxamide

Please amend the paragraph on page 81, lines 9-11, to read as follows:

Experiment 38

Preparation and Analysis of 116182

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(4-fluorophenylmethyl)carboxamide

Please amend the paragraph on page 81, lines 21-23, to read as follows:

Experiment 39

Preparation and Analysis of 116183

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(3,4-dimethoxyphenylmethyl)carboxamide

Please amend the paragraph on page 82, lines 1-3, to read as follows:

Experiment 40

Preparation and Analysis of 116184

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(3,4,5-trimethoxyphenylmethyl)carboxamide

Please amend the paragraph on page 82, lines 13-15, to read as follows:

Experiment 41

Preparation and Analysis of 116185

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-[(2-methylphenyl)methyl]amino]carboxamide

Please amend the paragraph on page 826, lines 25-27, to read as follows:

Experiment 42

Preparation and Analysis of 116189

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(4-methoxyphenylmethyl)carboxamide

Please amend the paragraph on page 83, lines 6-8, to read as follows:

Experiment 43


Preparation and Analysis of 116194

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-[2-(4-methoxy)phenylethyl]carboxamide

In the Claims:

Please delete claim 1. Please add new claims 14-26.

14. (New) A method for modulating by inverse agonism the activity of a human 5HT_{2A} serotonin receptor by contacting the receptor with a compound of formula:



Wherein:

W is lower alkyl (C₁₋₆), or halogen;

V is lower alkyl (C₁₋₆), or halogen;

X is either Oxygen or Sulfur;

Y is NR²R³, or (CH₂)_mR⁴, or O(CH₂)_nR⁴;

Z is lower alkyl (C₁₋₆);

m=0-4

n=0-4

R¹ is H or lower alkyl(C₁₋₄);

R² is H or lower alkyl(C₁₋₄);

R³ and R⁴ are independently a C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, OCF₃, SMe, COOR⁷, SO₂NR⁵R⁶, SO₃R⁷, CO-lower alkyl, SCF₃CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, and aryloxy wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₃R⁷, SO₂NR⁵R⁶, CO-lower alkyl, SCF₃CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl;

R^5 and R^6 are independently a H, or C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl, or CH_2 aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, $CONR^7R^8$, NR^7R^8 , $NHCOCH_3$, OCF_3 , SMe , $COOR^9$, SO_3R^7 , $SO_2NR^7R^8$, CO-lower alkyl, SCF_3 , CN, C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, and aryl wherein each of the C_{3-6} cycloalkyl, C_{1-6} alkyl, or aryl groups may be further optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, $CONR^8R^9$, NR^8R^9 , $NHCOCH_3$, OCF_3 , SMe , $COOR^7$, $SO_2NR^8R^9$, SO_3R^7 , CO-lower alkyl, SCF_3 , CN, C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, and aryl,


or R^5 and R^6 may form part of a 5, 6 or 7 membered cyclic structure which may be either saturated or unsaturated and that may contain up to four heteroatoms selected from O, N or S and said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , Me, NO_2 , OH, OMe, OEt, OCF_3 , SMe , $COOR^7$, $SO_2NR^8R^9$, SO_3R^7 , $HCOCH_3$, COEt, COMe, or halogen;

R^7 may be independently selected from H or C_{1-6} alkyl;

R^8 and R^9 are independently a H, or C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl, or CH_2 aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen, CF_3 , OCF_3 , OEt, CCl_3 , Me, NO_2 , OH, OMe, SMe , COMe, CN, $COOR^7$, SO_3R^7 , COEt, $NHCOCH_3$, or aryl;

an aryl moiety can be a 5 or 6 membered aromatic hetero-cyclic ring (containing up to 4 hetero atoms independently selected from N, O, or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle.

15. (New) A method for modulating by inverse agonism the activity of a human $5HT_{2A}$ serotonin receptor by contacting the receptor with a compound of formula:



(B)

Wherein:

W is Me, or Et, or halogen;

X is either Oxygen or Sulfur;

Y is NR^2R^3 , or $(\text{CH}_2)_m\text{R}^4$, or $\text{O}(\text{CH}_2)_n\text{R}^4$;

Z is lower alkyl (C_{1-6});

$m=0-4$

$n=0-4$

R^1 is H or lower alkyl (C_{1-4});

R^2 is H or lower alkyl (C_{1-4});

R^3 and R^4 are independently a C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH , CONR^5R^6 , NR^5R^6 , OCF_3 , SMe , COOR^7 , $\text{SO}_2\text{NR}^5\text{R}^6$, SO_3R^7 , CO-lower alkyl , SCF_3CN , C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, aryl, and aryloxy wherein each of the C_{3-6} cycloalkyl, C_{1-6} alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH , CONR^5R^6 , NR^5R^6 , NHCOCH_3 , OCF_3 , SMe , COOR^7 , SO_3R^7 , $\text{SO}_2\text{NR}^5\text{R}^6$, CO-lower alkyl , SCF_3CN , C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, and aryl;

R^5 and R^6 are independently a H, or C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl, or CH, aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, $CONR^7R^8$, NR^7R^8 , $NHCOCH_3$, OCF_3 , SMe, $COOR^9$, SO_3R^7 , $SO_2NR^7R^8$, CO-lower alkyl, SCF_3 , CN, C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, and aryl wherein each of the C_{3-6} cycloalkyl, C_{1-6} alkyl, or aryl groups may be further optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, $CONR^8R^9$, NR^8R^9 , $NHCOCH_3$, OCF_3 , SMe, $COOR^7$, $SO_2NR^8R^9$, SO_3R^7 , CO-lower alkyl, SCF_3 , CN, C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, and aryl,


or R^5 and R^6 may form part of a 5, 6 or 7 membered cyclic structure which may be either saturated or unsaturated and that may contain up to four heteroatoms selected from O, N or S and said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , Me, NO_2 , OH, OMe, OEt, OCF_3 , SMe, $COOR^7$, $SO_2NR^8R^9$, SO_3R^7 , $NHCOCH_3$, COEt, COMe, or halogen;

R^7 may be independently selected from H or C_{1-6} alkyl;

R^8 and R^9 are independently a H, or C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl, or CH_2 aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen, CF_3 , OCF_3 , OEt, CCl_3 , Me, NO_2 , OH, OMe, SMe, COMe, CN, $COOR^7$, SO_3R^7 , COEt, $NHCOCH_3$, or aryl;

an aryl moiety can be a 5 or 6 membered aromatic hetero-cyclic ring (containing up to 4 hetero atoms independently selected from N, O, or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle.

16. (New) A method for modulating by inverse agonism the activity of a human $5HT_{2A}$ serotonin receptor by contacting the receptor with a compound of formula:



Wherein:

R¹ and R² are H;

W is Br;

X is O;

Z is Me;

R³ is C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, OCF₃, SMe, COOR⁷, SO₂NR⁵R⁶, SO₃R⁷, CO-lower alkyl, SCF₃CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, and aryloxy wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₃R⁷, SO₂NR⁵R⁶, CO-lower alkyl, SCF₃CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl; R⁵ and R⁶ are independently a H, or C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl, or CH, aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁷R⁸, NR⁷R⁸, NHCOCH₃, OCF₃, SMe, COOR⁹, SO₃R⁷, SO₂NR⁷R⁸, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, or aryl groups may be further optionally substituted by up to four substituents in any position

independently selected from CF_3 , CCl_3 , NO_2 , OH , CONR^8R^9 , NR^8R^9 , NHCOCH_3 , OCF_3 , SMe , COOR^7 , $\text{SO}_2\text{NR}^8\text{R}^9$, SO_3R^7 , CO -lower alkyl, SCF_3 , CN , C_{2-6} alkenyl, H , halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, and aryl,


or R^5 and R^6 may form part of a 5, 6 or 7 membered cyclic structure which may be either saturated or unsaturated and that may contain up to four heteroatoms selected from O , N or S and said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , Me , NO_2 , OH , OMe , OEt , OCF_3 , SMe , COOR^7 , $\text{SO}_2\text{NR}^8\text{R}^9$, SO_3R^7 , NHCOCH_3 , COEt , COMe , or halogen;

R^7 may be independently selected from H or C_{1-6} alkyl;

R^8 and R^9 are independently a H , or C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl, or CH_2aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen, CF_3 , OCF_3 , OEt , CCl_3 , Me , NO_2 , OH , OMe , SMe , COMe , CN , COOR^7 , SO_3R^7 , COEt , NHCOCH_3 , or aryl;

an aryl moiety can be a 5 or 6 membered aromatic hetero-cyclic ring (containing up to 4 hetero atoms independently selected from N , O , or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle.

17. (New) A method for modulating by inverse agonism the activity of a human 5HT_{2A} serotonin receptor by contacting the receptor with a compound of formula:



Wherein:

W is Br;

X is O;

Z is Me;

R₁ is H

M = 0 – 4;

R⁴ is C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, OCF₃, SMe, COOR⁷, SO₂NR⁵R⁶, SO₃R⁷, CO-lower alkyl, SCF₃CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, and aryloxy wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₃R⁷, SO₂NR⁵R⁶, CO-lower alkyl, SCF₃CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl;

R⁵ and R⁶ are independently a H, or C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl, or CH, aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁷R⁸, NR⁷R⁸, NHCOCH₃, OCF₃, SMe, COOR⁹, SO₃R⁷, SO₂NR⁷R⁸, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens,

C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, or aryl groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁸R⁹, NR⁸R⁹, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₂NR⁸R⁹, SO₃R⁷, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl,


or R⁵ and R⁶ may form part of a 5, 6 or 7 membered cyclic structure which may be either saturated or unsaturated and that may contain up to four heteroatoms selected from O, N or S and said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, Me, NO₂, OH, OMe, OEt, OCF₃, SMe, COOR⁷, SO₂NR⁸R⁹, SO₃R⁷, NHCOCH₃, COEt, COMe, or halogen;

R⁷ may be independently selected from H or C₁₋₆ alkyl;

R⁸ and R⁹ are independently a H, or C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl, or CH₂aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen, CF₃, OCF₃, OEt, CCl₃, Me, NO₂, OH, OMe, SMe, COMe, CN, COOR⁷, SO₃R⁷, COEt, NHCOCH₃, or aryl;

an aryl moiety can be a 5 or 6 membered aromatic hetero-cyclic ring (containing up to 4 hetero atoms independently selected from N, O, or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle.

18. (New) A method for modulating by inverse agonism the activity of a human 5HT_{2A} serotonin receptor by contacting the receptor with a compound of formula:



wherein:

W is Br;

X is O;

Z is Me;

R¹ is H;

m = 0-4;

R⁴ is C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, OCF₃, SMe, COOR⁷, SO₂NR⁵R⁶, SO₃R⁷, CO-lower alkyl, SCF₃CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, and aryloxy wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₃R⁷, SO₂NR⁵R⁶, CO-lower alkyl, SCF₃CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl;

R⁵ and R⁶ are independently a H, or C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl, or CH, aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁷R⁸, NR⁷R⁸, NHCOCH₃, OCF₃, SMe, COOR⁹, SO₃R⁷, SO₂NR⁷R⁸, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, or

aryl groups may be further optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH , CONR^8R^9 , NR^8R^9 , NHCOCH_3 , OCF_3 , SMe , COOR^7 , $\text{SO}_2\text{NR}^8\text{R}^9$, SO_3R^7 , CO -lower alkyl, SCF_3 , CN , C_{2-6} alkenyl, H , halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, and aryl,





or R^5 and R^6 may form part of a 5, 6 or 7 membered cyclic structure which may be either saturated or unsaturated and that may contain up to four heteroatoms selected from O, N or S and said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , Me , NO_2 , OH , OMe , OEt , OCF_3 , SMe , COOR^7 , $\text{SO}_2\text{NR}^8\text{R}^9$, SO_3R^7 , NHCOCH_3 , COEt , COMe , or halogen;





R^7 may be independently selected from H or C_{1-6} alkyl;





R^8 and R^9 are independently a H , or C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl, or CH_2aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen, CF_3 , OCF_3 , OEt , CCl_3 , Me , NO_2 , OH , OMe , SMe , COMe , CN , COOR^7 , SO_3R^7 , COEt , NHCOCH_3 , or aryl;





an aryl moiety can be a 5 or 6 membered aromatic hetero-cyclic ring (containing up to 4 hetero atoms independently selected from N, O, or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle.





19. (New) The method of any one of claims 14-18 wherein the compound is selected from the group consisting of:














































, or



20. (New) A compound of formula (C):



(C)

Wherein:

W is Me, or Et, or halogen;

X is either Oxygen or Sulfur;

Y is NR^2R^3 , or $(\text{CH}_2)_m\text{R}^4$, or $\text{O}(\text{CH}_2)_n\text{R}^4$;

Z is lower alkyl (C_{1-6});

$m=0-4$;

$n=0-4$;

R^1 is H or lower alkyl (C_{1-4});

R^2 is H or lower alkyl (C_{1-4});

R^3 is a C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or $(\text{CH}_2)_k$ aryl group ($k=1-4$), and each said group may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, CONR^5R^6 , NR^5R^6 , OCF_3 , SMe, COOR^7 , $\text{SO}_2\text{NR}^5\text{R}^6$, SO_3R^7 , CO-lower alkyl, SCF_3 , CN, C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl,

aryl, and aryloxy wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₃R⁷, SO₂NR⁵R⁶, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl;

R⁴ is a C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, OCF₃, SMe, COOR⁷, SO₂NR⁵R⁶, SO₃R⁷, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, and aryloxy wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₃R⁷, SO₂NR⁵R⁶, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl;

R⁵ and R⁶ are independently a H, or C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl, or CH₂ aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁷R⁸, NR⁷R⁸, NHCOCH₃, OCF₃, SMe, COOR⁹, SO₃R⁷, SO₂NR⁷R⁸, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, or aryl groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁸R⁹, NR⁸R⁹, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₂NR⁸R⁹, SO₃R⁷, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl,

or R⁵ and R⁶ may form part of a 5, 6 or 7 membered cyclic structure which may be either saturated or unsaturated and that may contain up to four heteroatoms selected from O, N or S and said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, Me, NO₂, OH, OMe, OEt, OCF₃, SMe, COOR⁷,

$\text{SO}_2\text{NR}^8\text{R}^9$, SO_3R^7 , NHCOCH_3 , COEt , COMe , or halogen;

R^7 may be independently selected from H or C_{1-6} alkyl;

R^8 and R^9 are independently a H, or C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl, or CH_2 aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen, CF_3 , OCF_3 , OEt , CCl_3 , Me, NO_2 , OH, OMe, SMe, COMe , CN, COOR^7 , SO_3R^7 , COEt , NHCOCH_3 , or aryl;

an aryl moiety can be a 5 or 6 membered aromatic heterocyclic ring (containing up to 4 hetero atoms independently selected from N, O, or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle;


C_{1-6} alkyl moieties can be straight chain or branched;




optionally substituted C_{1-6} alkyl moieties can be straight chain or branched;





C_{2-6} alkenyl moieties can be straight chain or branched; and





optionally substituted C_{2-6} alkenyl moieties can be straight chain or branched.





21. (New) A compound having a structure selected from the group consisting of:





















































, or




22. (New) A compound structurally represented as follows:




23. (New) A compound structurally represented as follows:



24. (New) A compound structurally represented as follows:



25. (New) A compound structurally represented as follows:



- 26 (New) A composition comprising a compound of any one of claims 21-25.


REMARKS

Claim 1 was pending. Claim 1 has been cancelled without prejudice. New claims 14-26 have been added. No new matter has been added.

Upon entry of this amendment, claims 14-26 will be pending.

This is a continuation of U.S. Ser. No. 09/292,072, filed April 14, 1999, ("the 072 application"), which is presently pending. In this preliminary amendment, Applicants correct a nomenclature error in the parent application. This error arose from a mistaken structure assigned to a synthetic precursor. The synthetic precursor was purchased by Applicants from Maybridge


Chemical Company (Maybridge plc) under the designation KM 01978, and was said by Maybridge plc to be have the structure:



KM 01978

The compound was named by Maybridge plc 3-(4-bromo-1-methylpyrazole-3-yl)phenylamine. This was reported in the parent application at page 31, lines 2-7. Each of the compounds synthesized and characterized by Applicants, as reported in the 072 application in Example 13, Experiments 2-43, were prepared according to the synthetic procedures reported in the 072 application, using this starting material.

Recently, Applicants discovered that the compound provided to Applicants as KM 01978 in fact has the structure:



i.e., 3-(4-bromo-2-methylpyrazole-3-yl)phenylamine, which differs from the structure assigned by Maybridge plc in that the pyrazole methyl group is located on the adjacent nitrogen atom of the pyrazole ring. Accordingly, the name of each compound made and characterized by

Applicants, and deriving from the 3-(4-bromo-2-methylpyrazole-3-yl)phenylamine starting material, has been corrected. It is believed that no new matter is introduced by the correction, as the changes merely provide the accurate name of the compounds that Applicants prepared and characterized as described in the parent application. In addition, the generic formulae disclosed by Applicants also have been corrected. Inasmuch as those formulae were clearly based on the compounds prepared by Applicants in Example 13, Experiments 2-43, it is believed that the change to the generic structures also does not constitute new matter.

In addition, the specification has been amended to provide the correct structure and name of the Maybridge starting material, and compounds deriving therefrom, as discussed above.

Applicants present herein new claims 14 to 26. Claims 14 to 26 find support in the specification at, *inter alia*, page 31, line 8 to page 37, line 9; page 39; page 40; page 41; page 42; page 43; page 44, page 45; pages 47-49; page 51; pages 52-54; and pages 66-73.


Applicants attach hereto a paper and CRF version of the most recent Sequence Listing filed in the present application's parent.


Attached hereto is a marked-up version of the changes made to the specification and claims by the current amendment. The attached page is captioned "Version with markings to show changes made."

Applicants respectfully request that this amendment be entered without prejudice.

Applicants believe all of the claims presently before the Examiner patentably define the invention over the prior art and are otherwise in condition for ready allowance. Applicants respectfully request early notification of the same.

Respectfully submitted,


Gwilym J.O. Attwell
Registration No. 45,449


Michael P. Straher, Ph.D.
Registration No. 38,325

Date: *January 23, 2002*
WOODCOCK WASHBURN LLP
One Liberty Place - 46th Floor
Philadelphia, PA 19103
(215) 568-3100

VERSION WITH MARKINGS TO SHOW CHANGES MADE

In the Sequence Listing:

Please delete the Sequence Listing on file and insert therefore pages **1-19** comprising the most recently filed Sequence Listing.


In the Specification:

Please amend the paragraph on page 31, lines 2 to 7, as follows:


Based upon these results, structure activity analysis of the 103487 compound suggested that a series of derivatives of N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl] [(4-trifluoromethoxy)phenyl] aminocarboxamide [3-(4-bromo-1-methylpyrazole-3-yl)phenylamine] would exhibit similar 5-HT_{2A} activity and selectivity. A series of derivatives of N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl] [(4-trifluoromethoxy)phenyl] aminocarboxamide [3-(4-bromo-1-methylpyrazole-3-yl)phenylamine] were synthesized. These “directed” library compounds (Tripos, Inc.) were then analyzed in accordance with the protocols of Examples 9c(1), 9c(2) and 9d.

Please amend the paragraph on page 31, lines 8 to 12, as follows:

This series of compounds exhibits highly selective 5-HT_{2A} activity. Accordingly, in the first aspect of the invention, a series of compounds exhibiting 5-HT_{2A} receptor activity that are useful as inverse agonists at such receptors is designated by the general formula (A):




]




Please amend the paragraph on page 34, line 1, as follows:

[




]




Please amend the paragraph on page 36, lines 9-10, as follows:

[




]




Please amend the paragraph on page 37, lines 3-4, as follows:

[




]




Please amend the paragraph on page 37, lines 8-9, as follows:

[




]




Please amend the paragraph on page 39, lines 1-2, as follows:

[



]



Please amend the paragraphs spanning pages 39 to page 41 as follows:

Compound No.	R ¹	R ²	R ³	R ⁴	X	U	IP ₃ % of Control	IP ₃ AP-3 IC ₅₀ nM	WT 5HT _{2A} LSD IC ₅₀ nM
<u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]l[(4-methylthiophenyl)amino]carboxamide</u> <u>[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]l[(4-methylthiophenyl)amino]carboxamide]</u>									
116079	SCH ₃	H	H	H	O	NH	16	17	4
<u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]l[(4-chlorophenyl)amino]carboxamide</u> <u>[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]l[(4-chlorophenyl)amino]carboxamide]</u>									
116081	Cl	H	H	H	O	NH	10	3.2	11
<u>{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(4-fluorophenyl)carboxamide</u> <u>[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino}-N-(4-fluorophenyl)carboxamide]</u>									
116082	F	H	H	H	O	NH	11	-	7



<p><u>{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-[2-(trifluoromethoxy)phenyl]carboxamide</u> <u>[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino}-N-[2-(trifluoromethoxy)phenyl]carboxamide]</u></p>									
116087	H	H	CF ₃ O	H	O	NH	11	-	200
<p><u>{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(2-nitrophenyl)carboxamide</u> <u>[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino}-N-(2-nitrophenyl)carboxamide]</u></p>									
116089	H	H	NO ₂	H	O	NH	27	-	238
<p><u>{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(4-methoxyphenyl)carboxamide</u> <u>[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino}-N-(4-methoxyphenyl)carboxamide]</u></p>									
116091	MeO	H	H	H	O	NH	12	-	19
<p><u>{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(2-methylphenyl)carboxamide</u> <u>[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(2-methylphenyl)carboxamide]</u></p>									
116092	H	H	Me	H	O	NH	32	-	131
<p><u>{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-[4-(trifluoromethyl)phenyl]carboxamide</u> <u>[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino}-N-[4-(trifluoromethyl)phenyl]carboxamide]</u></p>									
116097	CF ₃	H	H	H	O	NH	11	-	65

<u>{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(3-chlorophenyl)carboxamide</u> <u>[[[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino]-N-(3-chlorophenyl)carboxamide]</u>									
116105	H	Cl	H	H	O	NH	11	-	39
<u>{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-[2-chlorophenyl]carboxamide</u> <u>[[[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino]-N-[2-chlorophenyl]carboxamide]</u>									
116108	H	H	Cl	H	O	NH	6	-	249
<u>{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-[4-(methylethyl)phenyl]carboxamide</u> <u>[[[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino]-N-[4-(methylethyl)phenyl]carboxamide]</u>									
116110	isopropyl	H	H	H	O	NH	7	-	338
<u>{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(3-methoxyphenyl)carboxamide</u> <u>[[[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino]-N-(3-methoxyphenyl)carboxamide]</u>									
116111	H	MeO	H	H	O	NH	7	-	106
<u>[[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino]-N-(3-methylphenyl)carboxamide</u> <u>[[[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino]-N-(3-methylphenyl)carboxamide]</u>									
116112	H	Me	H	H	O	NH	14	-	57

<p><u>[{3-(4-bromo-2-methylpyrazol-3-yl)phenyl}-amino]-N-methyl-N-[4-(trifluoromethoxy)phenyl]carboxamide</u></p> <p><u>[[{3-(4-bromo-1-methylpyrazol-3-yl)phenyl}-amino]-N-methyl-N-[4-(trifluoromethoxy)phenyl]carboxamide]</u></p>									
116113	CF ₃ O	H	H	H	O	NCH ₃	-	193	2
<p><u>N-[4-(tert-butyl)phenyl]{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}carboxamide</u></p> <p><u>[N-[4-(tert-butyl)phenyl]{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}carboxamide]</u></p>									
116119	t-butyl	H	H	H	O	NH	17	-	476
<p><u>N-[4-(dimethylamino)phenyl]{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}carboxamide</u></p> <p><u>[N-[4-(dimethylamino)phenyl]{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}carboxamide]</u></p>									
116122	NMe ₂	H	H	H	O	NH	9	-	309
<p><u>N-(3,5-dichloro-4-methylphenyl){[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}carboxamide</u></p> <p><u>[N-(3,5-dichloro-4-methylphenyl){[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}carboxamide]</u></p>									
116138	Me	Cl	H	Cl	O	NH	23	-	122

<u>{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-[4-(trifluoromethylthio)phenyl]carboxamide</u> <u>[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino}-N-[4-(trifluoromethylthio)phenyl]carboxamide]</u>									
116139	CF ₃ S	H	H	H	O	NH	12	-	56
<u>{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(2-fluorophenyl)carboxamide</u> <u>[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino}-N-(2-fluorophenyl)carboxamide]</u>									
116144	H	H	F	H	O	NH	12	-	37
<u>2-({[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}carbonylamino)benzamide</u> <u>[2-({[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino}carbonylamino)benzamide]</u>									
116145	H	H	CONH ₂	H	O	NH	31	-	7473
<u>{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(4-cyanophenyl)carboxamide</u> <u>[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino}-N-(4-cyanophenyl)carboxamide]</u>									
116147	CN	H	H	H	O	NH	12	-	2
<u>{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino}-N-(2-cyanophenyl)carboxamide</u> <u>[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino}-N-(2-cyanophenyl)carboxamide]</u>									
116148	H	H	CN	H	O	NH	30	-	348

[




Please amend the table on page 42 as follows:

Compound No.		IP ₃ AP-3	WT 5HT _{2A} LSD
		IC ₅₀ nM	IC ₅₀ nM


116141	<u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-</u> <u>[cyclohexylamino]carboxamide</u> [N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]- [cyclohexylamino]carboxamide]	114	81
--------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----	----

Please amend the paragraph on page 42, lines 3-4, as follows:

[



]




Please amend the table on page 42, lower, to 43, upper, as follows:


Compound No.	R ¹	R ²	R ³	R ⁴	R ⁵	IP ₃ AP-3	WT 5HT _{2A} LSD
						IC ₅₀ nM	IC ₅₀ nM
<u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-[phenylmethylamino]carboxamide</u> [N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-[phenylmethylamino]carboxamide]							
116143	H	H	H	H	H	120	47

<p><u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-[{(4-fluorophenyl)methyl} amino]carboxamide</u></p> <p>[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-[{(4-fluorophenyl)methyl} amino]carboxamide]</p>							
116182	F	H	H	H	H	89	132
<p><u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]- [{(3,4-dimethoxyphenyl)methyl} amino]-carboxamide</u></p> <p>[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]- [{(3,4-dimethoxyphenyl)methyl} amino]-carboxamide]</p>							
116183	OMe	OMe	H	H	H	-	1010
<p><u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-[{(3,4,5-trimethoxyphenyl)methyl} amino]-carboxamide</u></p> <p>[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-[{(3,4,5-trimethoxyphenyl)methyl} amino]-carboxamide]</p>							
116184	OMe	OMe	H	OMe	H	-	2960
<p><u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][{(2-methylphenyl)methyl} amino]carboxamide</u></p> <p>[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][{(2-methylphenyl)methyl} amino]carboxamide]</p>							
116185	H	H	Me	H	H	-	769

<p><u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][{(4-methoxyphenyl)methyl}amino]carboxamide</u></p> <p><u>[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][{(4-methoxyphenyl)methyl}amino]carboxamide]</u></p>							
116189	OMe	H	H	H	H	-	102

Please amend the paragraphs on pages 43-44, as follows:







Compound No.	R ¹	R ²	R ³	R ⁴	R ⁵	IP ₃ AP-3 IC ₅₀ nM	WT 5HT _{2A} LSD IC ₅₀ nM
N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][{2-(4-methoxyphenyl)ethyl}amino]carboxamide [N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][{2-(4-methoxyphenyl)ethyl}amino]carboxamide]							
116194	OMe	H	H	H	H	32	61

Please amend the paragraph on page 44, lines 7-8, as follows:

[





Please amend the paragraph on page 44, line 17, to page 45, line 2, as follows:

116100

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][4-methoxyphenoxy]carboxamide
[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][4-methoxyphenoxy]carboxamide]

[




Please amend the paragraph on page 45, lines 6-7 as follows:


116192

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(1,1-dimethylethoxy)carboxamide
 [{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(1,1-dimethylethoxy)carboxamide]

[





]



Please amend the paragraph on page 47, lines 3-4 as follows:

[



Please amend page 47, line 13, to page 49, line 1 as follows:


116101

$m = 0$, $R^1 = H$, $R^4 = 4\text{-trifluoromethoxyphenyl}$


[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][4-trifluoromethoxyphenyl]carboxamide]

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][4-trifluoromethoxyphenyl]carboxamide

[




]




116102

m=0, R¹= H, R⁴= thiopheneN-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][2-thienyl]carboxamide[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][2-thienyl]carboxamide]

[




]




116120

m=0, R¹= H, R⁴= chlorophenylN-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][4-chloro-phenyl]carboxamide[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][4-chloro-phenyl]carboxamide]

[





1



Please amend page 51, line 6, to page 52, line 1 as follows:

[




Name	Compound No.	R ¹	R ²	R ³	R ⁴	IP ₃	LSD
						IC ₅₀ nM	IC ₅₀ nM
[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-[4-(trifluoromethoxy)phenyl]acetamide]	116137	OCF ₃	H	H	H	-	106
<u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-[4-(trifluoromethoxy)phenyl]acetamide</u>							

[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-(3-fluorophenyl)acetamide] <u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(3-fluorophenyl)acetamide</u>	116174	H	F	H	H	153	318
[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-(3-methoxyphenyl)acetamide] <u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(3-methoxyphenyl)acetamide</u>	116175	H	OMe	H	H	108	625
[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-(2-fluorophenyl)acetamide] <u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(2-fluorophenyl)acetamide</u>	116176	H	H	F	H	129	662
[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-(4-nitrophenyl)acetamide] <u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(4-nitrophenyl)acetamide</u>	116177	NO ₂	H	H	H	61	108
<u>N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(2-methoxyphenyl)acetamide</u>	116178	H	H	OMe	H	165	2300


[compound names not provided]

Please amend the paragraph on page 52, line 7 as follows:

[




]




Please amend the paragraph on page 55, lines 5-13, as follows:


Compounds of general formula (I) can be obtained *via* a variety of synthetic routes all of which would be familiar to one skilled in the art. The reaction of isocyanates with amines is a commonly practiced method for the formation of ureas (see Org. Syn. Coll. Vol. V, (1973), 555). Amine (IV), 3-(4-bromo-2-methylpyrazole-3-yl)phenylamine [3-(4-bromo-1-methylpyrazole-3-yl)phenylamine], commercially available from Maybridge Chemical Company, Catalog No. KM01978, CAS No. 175201-77-1) reacts readily with isocyanates (V) in inert solvents such as halocarbons to yield the desired ureas of general formula (I) wherein $R^1 = R^2 = H$:




]



(IV)



(V)



(I) $\text{R}^1 = \text{R}^2 = \text{H}$


Please amend the paragraph spanning page 55, line 14, to page 56, line 1, as follows:

Alternatively the amine (IV) can be converted to the corresponding isocyanate (VI) by the action of phosgene or a suitable phosgene equivalent, *e.g.* triphosgene, in an inert

solvent such as a halocarbon in the presence of an organic base such as triethylamine or ethyldiisopropylamine. Isocyanate (VI) reacts with amines of general formula (VII), in an analogous fashion to that described above for the reaction of (IV) with (V), yielding the desired ureas of general formula (I) wherein $R^1 = H$:

[








Please amend the paragraph spanning page 56, line 2, to page 57, line 1, as follows:

Alternatively wherein the isocyanate of general formula (V) is not commercially available it can be prepared from the corresponding amine of general formula (VIII) in an analogous procedure to that described above for the preparation of (VI). Reaction of these isocyanates with (IV) would again yield the requisite ureas of general formula (I) wherein $R^1 = R^2 = H$:

[







Please amend the paragraph spanning page 57, line 2, to page 58, line 1, as follows:


Amines of general formula (VII) are also readily converted to activated isocyanate equivalents of general formula (IX) by the sequential action of carbonyldiimidazole and methyl iodide in tetrahydrofuran and acetonitrile respectively (R.A. Batey *et al*, *Tetrahedron Lett.*, (1998), 39, 6267-

6270.) Reaction of (IX) with (IV) in an inert solvent such as a halocarbon would yield the requisite ureas of general formula (I) wherein $R^1 = H$:

[

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
222








Please amend the paragraph spanning page 58, line 2, to page 59, line 1, as follows:

Amine (IV) may be monomethylated according to the procedure of J. Barluenga *et al*, *J. Chem. Soc., Chem. Commun.*, (1984), 20, 1334-1335, or alkylated according to the procedure of P. Marchini *et al*, *J. Org. Chem.*, (1975), 40(23), 3453-3456, to yield compounds of general formula (X) wherein $\text{R}^1 =$ lower alkyl. These materials may be reacted as above with reagents of general formula (V) and (IX) as



depicted below:







Please amend the paragraph on page 60, line 3, as follows:



[






Please amend the paragraph on page 60, line 4, as follows:

[




(II) $\text{R}^1 = \text{lower alkyl}$

]



(X) R¹ = lower alkyl

(XI)



(II) R¹ = lower alkyl

Please amend the paragraph on page 61, line 1, as follows:

[






]




Please amend the paragraph on page 62, line 1, as follows:

[




]




Please amend the paragraph on page 61, line 2, as follows:


[



(X) R^1 = lower alkyl





(XIII)




(III) R^1 = lower alkyl

]




Please amend page 63, line 1 as follows:

[




]




Please amend page 63, line 5, as follows:

[




(X) R¹ = lower alkyl




(XIV)

DCC/HOBT
or
HOBT/HBTU



(III) R¹ = lower alkyl

1



Please amend the paragraph on page 66, line 28 to page 67, line 2, as follows:

Experiment 2

Preparation and Analysis of 116100

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][4-methoxyphenoxy]carboxamide
 [N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][4-methoxyphenoxy]carboxamide]

To 4-methoxyphenylchloroformate (19 mg, 0.10 mmol) in CH_2Cl_2 (0.5 mL) was added dropwise a solution 3-(3-aminophenyl)-4-bromo-2-methylpyrazole [3-(3-aminophenyl)-4-bromo-1-methylpyrazole] (25 mg, 0.10 mmol) and triethylamine (14 μL , 0.10 mmol) in CH_2Cl_2 (0.5 mL). The mixture was stirred for 16 h and concentrated. Chromatography on flash silica (40% EtOAc/hexane) gave the title compound as a colourless solid (21 mg, 52%), m.p. 140.3-141.8°C. (EtOAc/hexane).

Please amend the paragraph on page 67, lines 9-17, as follows:

Experiment 3

Preparation and Analysis of 116101

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][4-trifluoromethoxyphenyl]carboxamide
[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][4-trifluoromethoxyphenyl]carboxamide]

To 4-(trifluoromethoxy)benzoyl chloride (19 μL , 0.12 mmol) in CH_2Cl_2 (1 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole [3-(3-aminophenyl)-4-bromo-1-methylpyrazole] (30 mg, 0.12 mmol) and triethylamine (17 μL , 0.12 mmol) in CH_2Cl_2 (0.5 mL). The reaction mixture was stirred for 16 h and concentrated. Chromatography on flash silica (50% EtOAc/hexane) gave the title compound as a colourless solid (40 mg, 76%), m.p. 138.6-139.6 °C (EtOAc/hexane).

Please amend the paragraph on page 67, lines 23-30, as follows:

Experiment 4

Preparation and Analysis of 116102

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][2-thienyl]carboxamide
[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][2-thienyl]carboxamide]

To thiophene-2-carbonyl chloride (11 μ L, 0.09 mmol) in CH_2Cl_2 (1 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole [3-(3-aminophenyl)-4-bromo-1-methylpyrazole] (25 mg, 0.09 mmol) and triethylamine (14 μ L, 0.09 mmol) in CH_2Cl_2 (0.5 mL). The reaction mixture was stirred for 16 h and concentrated. Chromatography on flash silica (50% EtOAc/hexane) gave the title compound as a colourless solid (24 mg, 68%), m.p. 127.8-128.6 $^\circ\text{C}$ (EtOAc/hexane).

Please amend the paragraph on page 68, lines 5-14, as follows:

Experiment 5

Preparation and Analysis of 116115

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][{(4-trifluoromethoxy)phenyl)methyl}amino]
carboxamide

[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][{(4-trifluoromethoxy)phenyl)
methyl}amino]carboxamide]

To a stirred solution of triphosgene (12 mg, 0.04 mmol) in CH_2Cl_2 (0.5 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole [3-(3-aminophenyl)-4-bromo-1-methylpyrazole] (30 mg, 0.12 mmol) and triethylamine (33 μ L, 0.24 mmol) in CH_2Cl_2 (0.5 mL). After 1 h, 4-(trifluoromethoxy)benzylamine (23 mg, 0.12 mmol) was added. The reaction mixture was stirred for 16 h and concentrated. Chromatography on flash silica (75% EtOAc/hexane) gave the title compound as a colourless solid (38 mg, 68%), m.p. 144.6-145.8 $^\circ\text{C}$ (EtOAc/hexane).

Please amend the paragraph on page 68, lines 21-28, as follows:

Experiment 6

Preparation and Analysis of 116120

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][4-chlorophenyl]carboxamide

[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][4-chlorophenyl]carboxamide]

To 4-chlorobenzoyl chloride (15 mg, 0.08 mmol) in CH₂Cl₂ (1 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole [3-(3-aminophenyl)-4-bromo-1-methylpyrazole] (21 mg, 0.08 mmol) and triethylamine (12 µL, 0.08 mmol) in CH₂Cl₂ (0.5 mL). The mixture was stirred for 16 h and concentrated. Chromatography on flash silica (50% EtOAc/hexane) gave the title compound as a colourless solid (23 mg, 72%), m.p. 184.4-184.8 °C (EtOAc/hexane).

Please amend the paragraph on page 69, lines 1-33, as follows:

Experiment 7

Preparation and Analysis of 116137

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-[4-(trifluoromethoxy)phenyl]acetamide
[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-[4-(trifluoromethoxy)phenyl]acetamide]

A solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole [3-(3-aminophenyl)-4-bromo-1-methylpyrazole] (35 mg, 0.14 mmol) and triethylamine (23 µL, 0.17 mmol) in DMF (0.5 mL) was added in one portion to a stirred solution of 4-trifluoromethoxyphenylacetic acid (31 mg, 0.14 mmol), HBTU (53 mg, 0.14 mmol) and HOBT (19 mg, 0.14 mmol) in DMF (1 mL). The mixture was heated at 70 °C for 24 h and then quenched with aqueous sodium bicarbonate solution. Ethyl acetate was added and the organic phase separated, washed with water (.times.3), brine, dried (MgSO₄) and evaporated. Chromatography on flash silica (50% EtOAc/hexane) gave the title compound as a colourless solid (43 mg, 68%). m.p. 141.2-142.5 °C (EtOAc/hexane).

Please amend the paragraph on page 69, lines 21-33, as follows:

Experiment 8

Preparation and Analysis of 116174

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(3-fluorophenyl)acetamide

[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-(3-fluorophenyl)acetamide]

A mixture of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole [3-(3-aminophenyl)-4-bromo-1-methylpyrazole] (30 mg, 0.12 mmol), 3-fluorophenylacetic acid (18 mg, 0.12 mmol), 1-hydroxybenzotriazole hydrate (16 mg, 0.12 mmol) and 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluoro-phosphate (46 mg, 0.12 mmol) were dissolved in chloroform (1.5 ml). N,N-Diisopropylethylamine (0.02 ml, 0.13 mmol) was added and the mixture stirred at room temperature for 16 h. The reaction mixture was then poured into brine and the organic layer washed with further brine, dried over magnesium sulphate and then concentrated in vacuo. The crude product was purified by column chromatography (ethyl acetate-toluene, 1:1), giving the title compound (12 mg, 26%). Rf 0.41 (ethyl acetate-toluene, 1:1).

Please amend the paragraph on page 70, lines 5-16, as follows:

Experiment 9

Preparation and Analysis of 116175

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(3-methoxyphenyl)acetamide

[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-(3-methoxyphenyl)acetamide]

A solution of 3-methoxyphenylacetyl chloride (0.02 ml, 0.12 mmol) in dichloromethane (0.75 ml) was added dropwise at 0 °C to a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole [3-(3-aminophenyl)-4-bromo-1-methylpyrazole] (30 mg, 0.12 mmol) and triethylamine (0.02 ml, 0.13 mmol) in dichloromethane (0.75 ml). The resulting mixture was stirred at room temperature for 16 h and then poured into brine. The organic layer was washed with more brine then dried over magnesium sulphate and concentrated in vacuo. The crude product was purified by column chromatography (ethyl acetate-toluene, 1:1), giving the title compound (9 mg, 19%). Rf 0.30 (ethyl acetate-toluene, 1:1).

Please amend the paragraph on page 71, lines 21-33, as follows:

Experiment 10

Preparation and Analysis of 116176

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(2-fluorophenyl)acetamide[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-(2-fluorophenyl)acetamide]

A mixture of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole [3-(3-aminophenyl)-4-bromo-1-methylpyrazole] (30 mg, 0.12 mmol), 2-fluorophenylacetic acid (18 mg, 0.12 mmol), 1-hydroxybenzotriazole hydrate (16 mg, 0.12 mmol) and 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluoro-phosphate (46 mg, 0.12 mmol) were dissolved in chloroform (1.5 ml). N,N-Diisopropylethylamine (0.02 ml, 0.13 mmol) was added and the mixture stirred at room temperature for 16 h. The reaction mixture was then poured into brine and the organic layer washed with further brine, dried over magnesium sulphate and then concentrated in vacuo. The crude product was purified by column chromatography (ethyl acetate-toluene, 1:1), giving the title compound (15 mg, 32%). Rf 0.52 (ethyl acetate-toluene, 1:1).

Please amend the paragraph on page 71, lines 5-17, as follows:

Experiment 11

Preparation and Analysis of 116177

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(4-nitrophenyl)acetamide[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-(4-nitrophenyl)acetamide]

A mixture of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole [3-(3-aminophenyl)-4-bromo-1-methylpyrazole] (30 mg, 0.12 mmol), 4-nitrophenylacetic acid (22 mg, 0.12 mmol), 1-hydroxybenzotriazole hydrate (16 mg, 0.12 mmol) and 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (46 mg, 0.12 mmol) were dissolved in chloroform (1.5 ml). N,N-Diisopropylethylamine (0.02 ml, 0.13 mmol) was added and the mixture stirred at room temperature for 16 h. The reaction mixture was then poured into brine and the organic layer washed with further brine, dried over magnesium sulphate and then concentrated in vacuo. The crude product was purified by column chromatography (ethyl acetate-toluene, 1:1), giving the title compound (9 mg, 18%). Rf 0.19 (ethyl acetate-toluene, 1:1).

Please amend the paragraph on page 71, line 22, to page 72, line 2, as follows:

Experiment 12

Preparation and Analysis of 116178

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-2-(2-methoxyphenyl)acetamide

[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-(2-methoxyphenyl)acetamide]

A mixture of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole [3-(3-aminophenyl)-4-bromo-1-methylpyrazole] (30 mg, 0.12 mmol), 2-methoxyphenylacetic acid (20 mg, 0.12 mmol), 1-hydroxybenzotriazole hydrate (16 mg, 0.12 mmol) and 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (46 mg, 0.12 mmol) were dissolved in chloroform (1.5 ml). N,N-Diisopropylethylamine (0.02 ml, 0.13 mmol) was added and the mixture stirred at room temperature for 16 h. The reaction mixture was then poured into brine and the organic layer washed with further brine, dried over magnesium sulphate and then concentrated in vacuo. The crude product was purified by column chromatography (chloroform-methanol, 99:1), giving the title compound (18 mg, 38%) as a colourless solid. R_f 0.65 (chloroform-methanol, 98:2).

Please amend the paragraph on page 72, lines 8-15, as follows:

Experiment 13

Preparation and Analysis of 116192

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(1,1-dimethylethoxy)carboxamide

[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(1,1-dimethylethoxy)carboxamide]

To di-tert-butyl dicarbonate (36 mg, 0.17 mmol) in methanol (1 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole [3-(3-aminophenyl)-4-bromo-1-methylpyrazole] (42 mg, 0.17 mmol) in methanol (1 mL). The mixture was stirred for 16 h and

concentrated. Chromatography on flash silica (40% EtOAc/hexane) gave the title compound as a colourless solid (29 mg, 49%) (EtOAc/hexane).

Please amend the paragraph on page 72, lines 22 to page 73, line 2, as follows:

One or the other (as indicated) of the two following synthetic protocols was used to generate each of the compounds below:

Protocol A:

To an isocyanate (1 mmol) in CH₂Cl₂ (4 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole [3-(3-aminophenyl)-4-bromo-1-methylpyrazole] (1 mmol) in CH₂Cl₂ (4 mL). The mixture was stirred for 16 hours and concentrated. Chromatography on flash silica (20%-80% EtOAc/hexane) followed by recrystallisation gave the pure urea.

Protocol B:

To a stirred solution of triphosgene (0.33 mmol) in CH₂Cl₂ (4 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-2-methylpyrazole [3-(3-aminophenyl)-4-bromo-1-methylpyrazole] (1 mmol) and triethylamine (2 mmol) in CH₂Cl₂ (4 mL). After 1 hour, an aniline was added (1 mmol). The reaction mixture was stirred for 16 hours and concentrated. Chromatography on flash silica (20%-80% EtOAc/hexane) followed by recrystallisation gave the pure urea.

Please amend the paragraph on page 73, lines 5-8, as follows:

Experiment 14

Preparation and Analysis of 116079

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][(4-methylthiophenyl)amino]carboxamide
[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][(4-methylthiophenyl)amino]carboxamide]

Please amend the paragraph on page 73, lines 16-18, as follows:

Experiment 15

Preparation and Analysis of 116081

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl][(4-chlorophenyl)amino]carboxamide
[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][(4-chlorophenyl)amino]carboxamide]

Please amend the paragraph on page 73, lines 29-31, as follows:

Experiment 16

Preparation and Analysis of 116082

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(4-fluorophenyl)carboxamide
[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(4-fluorophenyl)carboxamide]

Please amend the paragraph on page 74, lines 6-8, as follows:

Experiment 17

Preparation and Analysis of 116087

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-[2-(trifluoromethoxy)phenyl]carboxamide
[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-[2-(trifluoromethoxy)phenyl]carboxamide]

Please amend the paragraph on page 74, lines 17-19, as follows:

Experiment 18

Preparation and Analysis of 116089

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(2-nitrophenyl)carboxamide
[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(2-nitrophenyl)carboxamide]

Please amend the paragraph on page 74, lines 29-31, as follows:

Experiment 19

Preparation and Analysis of 116091

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(4-methoxyphenyl)carboxamide
[{{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(4-methoxyphenyl)carboxamide]

Please amend the paragraph on page 75, lines 7-9, as follows:

Experiment 20

Preparation and Analysis of 116092

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(2-methylphenyl)carboxamide
[{{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(2-methylphenyl)carboxamide]

Please amend the paragraph on page 75, lines 18-20, as follows:

Experiment 21

Preparation and Analysis of 116097

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-[4-(trifluoromethyl)phenyl]carboxamide
[{{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-[4-(trifluoromethyl)phenyl]carboxamide]

Please amend the paragraph on page 75, lines 28-30, as follows:

Experiment 22

Preparation and Analysis of 116105

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(3-chlorophenyl)carboxamide

[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(3-chlorophenyl)carboxamide]

Please amend the paragraph on page 76, lines 8-10, as follows:

Experiment 23

Preparation and Analysis of 116108

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(2-chlorophenyl)carboxamide

[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(2-chlorophenyl)carboxamide]

Please amend the paragraph on page 76, lines 20-22, as follows:

Experiment 24

Preparation and Analysis of 116110

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-[4-(methylethyl)phenyl]carboxamide

[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-[4-(methylethyl)phenyl]carboxamide]

Please amend the paragraph on page 76, lines 31-33, as follows:

Experiment 25

Preparation and Analysis of 116111

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(3-methoxyphenyl)carboxamide

[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(3-methoxyphenyl)carboxamide]

Please amend the paragraph on page 77, lines 8-10, as follows:

Experiment 26

Preparation and Analysis of 116112

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(3-methylphenyl)carboxamide

[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(3-methylphenyl)carboxamide]

Please amend the paragraph on page 77, lines 19-22, as follows:

Experiment 27

Preparation and Analysis of 116113

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-methyl-N-[4-(trifluoromethoxy)phenyl]-carboxamide

[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-methyl-N-[4-(trifluoromethoxy)phenyl]-carboxamide]

Please amend the paragraph on page 77, lines 30-32, as follows:

Experiment 28

Preparation and Analysis of 116119

N-[4-(tert-butyl)phenyl]{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}carboxamide

[N-[4-(tert-butyl)phenyl]{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}carboxamide]

Please amend the paragraph on page 78, lines 8-10, as follows:

Experiment 29

Preparation and Analysis of 116122

N-[4-(dimethylamino)phenyl]{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}carboxamide

[N-[4-(dimethylamino)phenyl]{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}carboxamide]

Please amend the paragraph on page 78, lines 19-21, as follows:

Experiment 30

Preparation and Analysis of 116138

N-(3,5-dichloro-4-methylphenyl){[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}carboxamide

[N-[4-(dimethylamino)phenyl]{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}carboxamide]

Please amend the paragraph on page 78, lines 30-32, as follows:

Experiment 31

Preparation and Analysis of 116139

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-[4-(trifluoromethylthio)phenyl]carboxamide
[[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino]-N-[4-(trifluoromethylthio)phenyl]carboxamide]

Please amend the paragraph on page 79, lines 8-10, as follows:

Experiment 32

Preparation and Analysis of 116141

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(cyclohexyl)carboxamide
[[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino]-N-(cyclohexyl)carboxamide]

Please amend the paragraph on page 79, lines 21-23, as follows:

Experiment 33

Preparation and Analysis of 116143

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(phenylmethyl)carboxamide
[[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino]-N-(phenylmethyl)carboxamide]

Please amend the paragraph on page 80, lines 1-3, as follows:

Experiment 34

Preparation and Analysis of 116144

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(2-fluorophenyl)carboxamide
[[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino]-N-(2-fluorophenyl)carboxamide]

Please amend the paragraph on page 80, lines 11-13, as follows:

Experiment 35

Preparation and Analysis of 116145

2-([3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-amino)carbonylamino)benzamide

[2-([3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-amino)carbonylamino)benzamide]

Please amend the paragraph on page 80, lines 21-23, as follows:

Experiment 36

Preparation and Analysis of 116147

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(4-cyanophenyl)carboxamide

[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(4-cyanophenyl)carboxamide]

Please amend the paragraph on page 80, lines 31-33, as follows:

Experiment 37

Preparation and Analysis of 116148

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(2-cyanophenyl)carboxamide

[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(2-cyanophenyl)carboxamide]

Please amend the paragraph on page 81, lines 9-11, as follows:

Experiment 38

Preparation and Analysis of 116182

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(4-fluorophenylmethyl)carboxamide

[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(4-fluorophenylmethyl)carboxamide]

Please amend the paragraph on page 81, lines 21-23, as follows:

Experiment 39

Preparation and Analysis of 116183

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(3,4-
dimethoxyphenylmethyl)carboxamide
[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(3,4-
dimethoxyphenylmethyl)carboxamide]

Please amend the paragraph on page 82, lines 1-3, as follows:

Experiment 40

Preparation and Analysis of 116184

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(3,4,5-
trimethoxyphenylmethyl)carboxamide
[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(3,4,5-
trimethoxyphenylmethyl)carboxamide]

Please amend the paragraph on page 82, lines 13-15, as follows:

Experiment 41

Preparation and Analysis of 116185

N-[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]-[{(2-methylphenyl)methyl } amino]carboxamide
[N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-[{(2-methylphenyl)methyl } amino]carboxamide]

Please amend the paragraph on page 826, lines 25-27, as follows:

Experiment 42

Preparation and Analysis of 116189

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-(4-methoxyphenylmethyl)carboxamide[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(4-methoxyphenylmethyl)carboxamide]

Please amend the paragraph on page 83, lines 6-8, as follows:


Experiment 43

Preparation and Analysis of 116194

{[3-(4-bromo-2-methylpyrazol-3-yl)phenyl]amino}-N-[2-(4-methoxy)phenylethyl]carboxamide[{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-[2-(4-methoxy)phenylethyl]carboxamide]**In the Claims:**

Please delete claim 1. Please add new claims 14-26.

14. (New) A method for modulating by inverse agonism the activity of a human 5HT_{2A} serotonin receptor by contacting the receptor with a compound of formula:



Wherein:

W is lower alkyl (C₁₋₆), or halogen;

V is lower alkyl (C₁₋₆), or halogen;

X is either Oxygen or Sulfur;

Y is NR²R³, or (CH₂)_mR⁴, or O(CH₂)_nR⁴;

Z is lower alkyl (C₁₋₆);

m=0-4

n=0-4

R¹ is H or lower alkyl(C₁₋₄);

R² is H or lower alkyl(C₁₋₄);

R³ and R⁴ are independently a C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, OCF₃, SMe, COOR⁷, SO₂NR⁵R⁶, SO₃R⁷, CO-lower alkyl, SCF₃CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, and aryloxy wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₃R⁷, SO₂NR⁵R⁶, CO-lower alkyl, SCF₃CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl;

R⁵ and R⁶ are independently a H, or C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl, or CH₂ aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁷R⁸, NR⁷R⁸, NHCOCH₃, OCF₃, SMe, COOR⁹, SO₃R⁷, SO₂NR⁷R⁸, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, or aryl groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁸R⁹, NR⁸R⁹, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₂NR⁸R⁹, SO₃R⁷, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl,

or R⁵ and R⁶ may form part of a 5, 6 or 7 membered cyclic structure which may be either


saturated or unsaturated and that may contain up to four heteroatoms selected from O, N or S and said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, Me, NO₂, OH, OMe, OEt, OCF₃, SMe, COOR⁷, SO₂NR⁸R⁹, SO₃R⁷, HCOCH₃, COEt, COMe, or halogen;

R⁷ may be independently selected from H or C₁₋₆ alkyl;

R⁸ and R⁹ are independently a H, or C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl, or CH₂ aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen, CF₃, OCF₃, OEt, CCl₃, Me, NO₂, OH, OMe, SMe, COMe, CN, COOR⁷, SO₃R⁷, COEt, NHCOCH₃, or aryl;

an aryl moiety can be a 5 or 6 membered aromatic hetero-cyclic ring (containing up to 4 hetero atoms independently selected from N, O, or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle.

15. (New) A method for modulating by inverse agonism the activity of a human 5HT_{2A} serotonin receptor by contacting the receptor with a compound of formula:



(B)

Wherein:

W is Me, or Et, or halogen;

X is either Oxygen or Sulfur;

Y is NR^2R^3 , or $(\text{CH}_2)_m\text{R}^4$, or $\text{O}(\text{CH}_2)_n\text{R}^4$;

Z is lower alkyl (C_{1-6});

$m=0-4$

$n=0-4$

R^1 is H or lower alkyl (C_{1-4});

R^2 is H or lower alkyl (C_{1-4});

R^3 and R^4 are independently a C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, CONR^5R^6 , NR^5R^6 , OCF_3 , SMe, COOR^7 , $\text{SO}_2\text{NR}^5\text{R}^6$, SO_3R^7 , CO-lower alkyl, SCF_3CN , C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, aryl, and aryloxy wherein each of the C_{3-6} cycloalkyl, C_{1-6} alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, CONR^5R^6 , NR^5R^6 , NHCOCH_3 , OCF_3 , SMe, COOR^7 , SO_3R^7 , $\text{SO}_2\text{NR}^5\text{R}^6$, CO-lower alkyl, SCF_3CN , C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, and aryl;

R^5 and R^6 are independently a H, or C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl, or CH, aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, CONR^7R^8 , NR^7R^8 , NHCOCH_3 , OCF_3 , SMe, COOR^9 , SO_3R^7 , $\text{SO}_2\text{NR}^7\text{R}^8$, CO-lower alkyl, SCF_3 , CN, C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, and aryl wherein each of the C_{3-6} cycloalkyl, C_{1-6} alkyl, or aryl groups may be further optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, CONR^8R^9 , NR^8R^9 , NHCOCH_3 , OCF_3 , SMe, COOR^7 , $\text{SO}_2\text{NR}^8\text{R}^9$, SO_3R^7 , CO-lower alkyl, SCF_3 , CN, C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, and aryl,

or R^5 and R^6 may form part of a 5, 6 or 7 membered cyclic structure which may be either saturated or unsaturated and that may contain up to four heteroatoms selected from O, N or S and


said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , Me, NO_2 , OH, OMe, OEt, OCF_3 , SMe, COOR^7 , $\text{SO}_2\text{NR}^8\text{R}^9$, SO_3R^7 , NHCOCH_3 , COEt, COMe, or halogen;

R^7 may be independently selected from H or C_{1-6} alkyl;

R^8 and R^9 are independently a H, or C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl, or CH_2aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen, CF_3 , OCF_3 , OEt, CCl_3 , Me, NO_2 , OH, OMe, SMe, COMe, CN, COOR^7 , SO_3R^7 , COEt, NHCOCH_3 , or aryl;

an aryl moiety can be a 5 or 6 membered aromatic hetero-cyclic ring (containing up to 4 hetero atoms independently selected from N, O, or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle.

16. (New) A method for modulating by inverse agonism the activity of a human 5HT_{2A} serotonin receptor by contacting the receptor with a compound of formula:



Wherein:

R¹ and R² are H;

W is Br;

X is O;

Z is Me;

R^3 is C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, $CONR^5R^6$, NR^5R^6 , OCF_3 , SMe, $COOR^7$, $SO_2NR^5R^6$, SO_3R^7 , CO-lower alkyl, SCF_3CN , C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, aryl, and aryloxy wherein each of the C_{3-6} cycloalkyl, C_{1-6} alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, $CONR^5R^6$, NR^5R^6 , $NHCOCH_3$, OCF_3 , SMe, $COOR^7$, SO_3R^7 , $SO_2NR^5R^6$, CO-lower alkyl, SCF_3CN , C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, and aryl; R^5 and R^6 are independently a H, or C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl, or CH, aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, $CONR^7R^8$, NR^7R^8 , $NHCOCH_3$, OCF_3 , SMe, $COOR^9$, SO_3R^7 , $SO_2NR^7R^8$, CO-lower alkyl, SCF_3 , CN, C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, and aryl wherein each of the C_{3-6} cycloalkyl, C_{1-6} alkyl, or aryl groups may be further optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, $CONR^8R^9$, NR^8R^9 , $NHCOCH_3$, OCF_3 , SMe, $COOR^7$, $SO_2NR^8R^9$, SO_3R^7 , CO-lower alkyl, SCF_3 , CN, C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, and aryl,

or R^5 and R^6 may form part of a 5, 6 or 7 membered cyclic structure which may be either saturated or unsaturated and that may contain up to four heteroatoms selected from O, N or S and said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , Me, NO_2 , OH, OMe, OEt, OCF_3 , SMe, $COOR^7$, $SO_2NR^8R^9$, SO_3R^7 , $NHCOCH_3$, COEt, COMe, or halogen;


R^7 may be independently selected from H or C_{1-6} alkyl;

R^8 and R^9 are independently a H, or C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl, or CH_2 aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen, CF_3 , OCF_3 , OEt, CCl_3 ,

Me, NO₂, OH, OMe, SMe, COMe, CN, COOR⁷, SO₃R⁷, COEt, NHCOCH₃, or aryl;

an aryl moiety can be a 5 or 6 membered aromatic hetero-cyclic ring (containing up to 4 hetero atoms independently selected from N, O, or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle.

17. (New) A method for modulating by inverse agonism the activity of a human 5HT_{2A} serotonin receptor by contacting the receptor with a compound of formula:



Wherein:

W is Br;

X is O;

Z is Me;

R¹ is H

n = 0 - 4;

R⁴ is C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, OCF₃, SMe, COOR⁷, SO₂NR⁵R⁶, SO₃R⁷, CO-lower alkyl,

SCF₃CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, and aryloxy wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₃R⁷, SO₂NR⁵R⁶, CO-lower alkyl, SCF₃CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl;

R⁵ and R⁶ are independently a H, or C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl, or CH, aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁷R⁸, NR⁷R⁸, NHCOCH₃, OCF₃, SMe, COOR⁹, SO₃R⁷, SO₂NR⁷R⁸, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, or aryl groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁸R⁹, NR⁸R⁹, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₂NR⁸R⁹, SO₃R⁷, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl,


or R⁵ and R⁶ may form part of a 5, 6 or 7 membered cyclic structure which may be either saturated or unsaturated and that may contain up to four heteroatoms selected from O, N or S and said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, Me, NO₂, OH, OMe, OEt, OCF₃, SMe, COOR⁷, SO₂NR⁸R⁹, SO₃R⁷, NHCOCH₃, COEt, COMe, or halogen;

R⁷ may be independently selected from H or C₁₋₆ alkyl;

R⁸ and R⁹ are independently a H, or C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl, or CH₂aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen, CF₃, OCF₃, OEt, CCl₃, Me, NO₂, OH, OMe, SMe, COMe, CN, COOR⁷, SO₃R⁷, COEt, NHCOCH₃, or aryl;

an aryl moiety can be a 5 or 6 membered aromatic hetero-cyclic ring (containing up to 4 hetero atoms independently selected from N, O, or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle.

18. (New) A method for modulating by inverse agonism the activity of a human $5HT_{2A}$ serotonin receptor by contacting the receptor with a compound of formula:



wherein:

W is Br;

X is O;

Z is Me;

R1 is H;

m= 0-4;

R^4 is C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, $CONR^5R^6$, NR^5R^6 , OCF_3 , SMe, $COOR^7$, $SO_2NR^5R^6$, SO_3R^7 , CO-lower alkyl, SCF_3CN , C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, aryl, and aryloxy wherein each of the C_{3-6} cycloalkyl, C_{1-6} alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, $CONR^5R^6$, NR^5R^6 , $NHCOCH_3$, OCF_3 , SMe, $COOR^7$, SO_3R^7 , $SO_2NR^5R^6$, CO-lower alkyl, SCF_3CN , C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl, and aryl;

R^5 and R^6 are independently a H, or C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or aryl, or

CH, aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁷R⁸, NR⁷R⁸, NHCOCH₃, OCF₃, SMe, COOR⁹, SO₃R⁷, SO₂NR⁷R⁸, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, or aryl groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁸R⁹, NR⁸R⁹, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₂NR⁸R⁹, SO₃R⁷, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl,




or R⁵ and R⁶ may form part of a 5, 6 or 7 membered cyclic structure which may be either saturated or unsaturated and that may contain up to four heteroatoms selected from O, N or S and said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, Me, NO₂, OH, OMe, OEt, OCF₃, SMe, COOR⁷, SO₂NR⁸R⁹, SO₃R⁷, NHCOCH₃, COEt, COMe, or halogen;





R⁷ may be independently selected from H or C₁₋₆ alkyl;





R⁸ and R⁹ are independently a H, or C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl, or CH₂aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen, CF₃, OCF₃, OEt, CCl₃, Me, NO₂, OH, OMe, SMe, COMe, CN, COOR⁷, SO₃R⁷, COEt, NHCOCH₃, or aryl;





an aryl moiety can be a 5 or 6 membered aromatic hetero-cyclic ring (containing up to 4 hetero atoms independently selected from N, O, or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle.





19. (New) The method of any one of claims 14-18 wherein the compound is selected from the group consisting of:


















































, or



20. (New) A compound of formula (C):



(C)

Wherein:

W is Me, or Et, or halogen;

X is either Oxygen or Sulfur;

Y is NR^2R^3 , or $(\text{CH}_2)_m\text{R}^4$, or $\text{O}(\text{CH}_2)_n\text{R}^4$;

Z is lower alkyl (C_{1-6});

$m=0-4$;

$n=0-4$;

R^1 is H or lower alkyl (C_{1-4});

R^2 is H or lower alkyl (C_{1-4});

R^3 is a C_{1-6} alkyl, or C_{2-6} alkenyl, or cycloalkyl, or $(\text{CH}_2)_k$ aryl group ($k=1-4$), and each said group may be optionally substituted by up to four substituents in any position independently selected from CF_3 , CCl_3 , NO_2 , OH, CONR^5R^6 , NR^5R^6 , OCF_3 , SMe , COOR^7 , $\text{SO}_2\text{NR}^5\text{R}^6$, SO_3R^7 , CO-lower alkyl, SCF_3 CN, C_{2-6} alkenyl, H, halogens, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkyl,

aryl, and aryloxy wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₃R⁷, SO₂NR⁵R⁶, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl;

R⁴ is a C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, OCF₃, SMe, COOR⁷, SO₂NR⁵R⁶, SO₃R⁷, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, and aryloxy wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁵R⁶, NR⁵R⁶, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₃R⁷, SO₂NR⁵R⁶, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl;

R⁵ and R⁶ are independently a H, or C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl, or CH₂ aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁷R⁸, NR⁷R⁸, NHCOCH₃, OCF₃, SMe, COOR⁹, SO₃R⁷, SO₂NR⁷R⁸, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl wherein each of the C₃₋₆ cycloalkyl, C₁₋₆ alkyl, or aryl groups may be further optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, NO₂, OH, CONR⁸R⁹, NR⁸R⁹, NHCOCH₃, OCF₃, SMe, COOR⁷, SO₂NR⁸R⁹, SO₃R⁷, CO-lower alkyl, SCF₃, CN, C₂₋₆ alkenyl, H, halogens, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, C₁₋₆ alkyl, and aryl,

or R⁵ and R⁶ may form part of a 5, 6 or 7 membered cyclic structure which may be either saturated or unsaturated and that may contain up to four heteroatoms selected from O, N or S and said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from CF₃, CCl₃, Me, NO₂, OH, OMe, OEt, OCF₃, SMe, COOR⁷,

$\text{SO}_2\text{NR}^8\text{R}^9$, SO_3R^7 , NHCCH_3 , COEt , COMe , or halogen;

R⁷ may be independently selected from H or C₁₋₆ alkyl;

R⁸ and R⁹ are independently a H, or C₁₋₆ alkyl, or C₂₋₆ alkenyl, or cycloalkyl, or aryl, or CH₂ aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen, CF₃, OCF₃, OEt, CCl₃, Me, NO₂, OH, OMe, SMe, COMe, CN, COOR⁷, SO₃R⁷, COEt, NHCOCH₃, or aryl;

an aryl moiety can be a 5 or 6 membered aromatic heterocyclic ring (containing up to 4 hetero atoms independently selected from N, O, or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle;


C₁₋₆ alkyl moieties can be straight chain or branched;




optionally substituted C₁₋₆ alkyl moieties can be straight chain or branched;





C₂₋₆ alkenyl moieties can be straight chain or branched; and





optionally substituted C₂₋₆ alkenyl moieties can be straight chain or branched.





21. **(New)** A compound having a structure selected from the group consisting of:





















































, or




22. (New) A compound structurally represented as follows:




23. (New) A compound structurally represented as follows:



24. (New) A compound structurally represented as follows:



25. (New) A compound structurally represented as follows:



- 26 (New) A composition comprising a compound of any one of claims 21-25.